

Erratum - Long-term follow-up of trans-obturator vaginal tape for stress urinary incontinence: a single-center retrospective study

Prisca Sozzani, Carola Minella, Pier Luigi Montironi, Francesca Longo, Andrea Scoletta

The following Table is the correct version of Table 1 published in Urogynaecologia 2026; 38:368 (doi:10.4081/uij.2026.368).

Table 1. Baseline characteristics of patients included in the study.

Characteristics	Group 1 (<5 years follow-up)	Group 2 (≥5 years follow-up)	p
Age at surgery, years, median (range)	57 (37-77)	57 (46-75)	0.93
BMI, kg/m ² , mean ± standard deviation	25.37±3.94	27.11±3.52	0.12
Parity, median (range)	2 (0-5)	2 (0-7)	0.81
Fetal macrosomia, n (%)	4 (16.7)	5 (26.3)	0.44
Smoke, n (%)	6 (25)	6 (31.6)	0.63
Previous gynecological surgery, n (%)	3 (12.5)	5 (26.3)	0.25
Anterior prolapse, grade Baden-Walker 1, n (%)	8 (33.3)	9 (47.4)	0.35
MPCU, mean ± standard deviation	56.58±20.36	60.89±23.71	0.52
Detrusor overactivity, n (%)	3 (12.5)	2 (10.5)	0.84
Normal compliance, n (%)	24 (100)	19 (100)	/
Type of incontinence			0.99
SUI, n (%)	19 (79.2)	15 (78.9)	
MUI, n (%)	5 (20.8)	4 (21.1)	
Follow-up, years, median (range)	4 (1-5)	11 (7-13)	/

BMI, body mass index; MPCU, maximum urethral closure pressure; SUI, stress urinary incontinence; MUI, mixed urinary incontinence.